**Characteristics of 1,719 participants from the Cancer 2015 study who provided EORTC QLQ-C30 data (Woodcock and Doble, 2018)**

| Characteristic | % |
|---|---|
| % female | 45.4 |
| Age (Mean, standard deviation) | 61.6, 12.5 |
| Recruiting hospital | |
|     Public | 81.2 |
|     Private | 16.8 |
| Smoking status | |
|     Current smoker | 14.8 |
|     Ex-smoker | 46.2 |
|     Never smoked | 39.0 |
| ECOG score | |
|     0, Normal activity | 69.8 |
|     1, Limited | 22.1 |
|     2, Self-care | 6.3 |
|     3, Limited self-care | 1.9 |
|     4, No self-care | 0.1 |
| Tumour site | |
|     Breast | 20.2 |
|     Prostate | 16.2 |
|     Head and neck | 15.2 |
|     Colorectal | 10.4 |
|     Lung | 8.6 |
|     Oesophagogastric | 3.4 |
|     Renal | 4.4 |
|     Bone and soft tissue | 3.2 |
|     Cancer of unknown primary | 1.7 |
|     Cervical | 2.3 |
|     Other | 14.4 |
| Stage | |
|     0 | 1.0 |
|     1 | 26.2 |
|     2 | 29.5 |
|     3 | 21.4 |
|     4 | 19.5 |
|     Other | 2.5 |

**Reference**

Woodcock, F., & Doble, B. (2018). Mapping the EORTC-QLQ-C30 to the EQ-5D-3L: an assessment of existing and newly developed algorithms. Medical Decision Making, 38(8), 954-967.